

Where is the π resonance?

O. Tchernyshyov,^{1,2} M. R. Norman,² and A. V. Chubukov^{2,3}

¹*School of Natural Sciences, Institute for Advanced Study, Princeton, New Jersey, 08540*

²*Materials Science Division, Argonne National Laboratory, Argonne, Illinois 60439*

³*Department of Physics, University of Wisconsin, Madison, Wisconsin 53706*

(September 5, 2000)

We discuss the interplay of particle-particle and particle-hole spin-triplet channels in high- T_c superconductors using a quasiparticle dispersion motivated by angle-resolved photoemission. Within a generalized RPA, we find a well defined antibound state of two holes, the π resonance of Demler and Zhang, as well as a bound state of a particle and a hole, the spin exciton. We show that the energy of the π resonance always exceeds 2Δ , twice the maximum d-wave gap, therefore the neutron resonance observed in the cuprates around energy Δ is most likely a spin exciton. At the same time, we speculate that the π particle can exist at higher energies and might be observed in neutron scattering around 100 meV.

74.20.-z, 74.72.-h, 76.50.+g

Reflecting our poor understanding of high- T_c superconductivity in general, theoretical debates continue on virtually every aspect of it. A good example is the resonance observed in inelastic neutron scattering at energies 25–43 meV in $\text{YBa}_2\text{Cu}_3\text{O}_{7-y}$ and $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$.^{1,2} Measurements using polarized neutrons³ indicate that the observed excitation is magnetic in nature. Theoretical proposals include a van Hove singularity in the Stoner continuum,^{4,5} a bound state (spin exciton) in the particle-hole channel,^{6–10} and an antibound state in the particle-particle channel (π resonance).^{11,12} Deciding between these approaches is somewhat difficult since in all theories the resonance mode is an excited pair of quasiparticles with total lattice momentum $\mathbf{Q} = (\pi/a, \pi/a)$ and spin $S = 1$.^{9,12,13}

In this paper, we use available experimental data and a simple kinematic argument to demonstrate that a particle-hole bound state is the most likely explanation for the resonance. In a nutshell, our reasoning goes as follows. Angle-resolved photoemission spectroscopy (ARPES) directly measures the dispersion of fermionic quasiparticles $\epsilon_{\mathbf{k}}$ in the superconducting state. From this, one can deduce information about the continuum of states of two quasiparticles with total momentum $\mathbf{Q}$ and, in particular, determine its energy bounds. An analysis of ARPES data in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ shows that the two-quasiparticle continuum starts at $E_{\min} \approx 2\Delta$, where Δ is the maximum value of the superconducting gap at the Fermi surface. The commensurate neutron resonance in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ resides at $E_{\text{res}} \approx \Delta$, definitely below the lower edge of the continuum. The resonance then can only be a bound state of two quasiparticles, not an antibound state.

This argument—that the lower edge of the two-particle continuum is determined by Δ and not by the chemical potential μ as in Ref. 11—is based on the analysis of the Fermi surface inferred from ARPES data (Fig. 1). We have verified that, for this Fermi surface, $E_{\min}$ is determined not by electrons along zone diagonals $k_x = k_y$, but rather by electrons near the hot spots (points at $\mathbf{k} = \mathbf{k}_F$

separated by $\mathbf{Q}$). The latter are located close to $(0, \pi)$ and symmetry related points, and in the superconducting state have a gap $\Delta(\mathbf{k}_{\text{hs}}) \approx \Delta$. This yields a threshold at $E_{\min} \approx 2\Delta$. Note that this argument also invalidates the interpretation of the resonance as a van Hove singularity in the two-quasiparticle continuum.⁵

Does it mean that the π resonance does not exist? Not necessarily. In our model calculations, the spin exciton and the π resonance are found to coexist at intermediate coupling strengths, similar to earlier findings.¹² Furthermore, in a situation where the continuum of states with two holes is narrow and has a sharp upper edge, we find that the π resonance is rather sharp. To which extent these conditions are satisfied in the cuprates is not clear because self-energy effects could wash out the upper edge of the continuum. We therefore can only speculate that the π resonance might be discovered in neutron scattering at energies above $E_{\min}$ (60 meV in optimally doped $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$), more likely around 100 meV.

To proceed, we need a model treating all spin-triplet excitations (with charges 0 and ± 2) on an equal footing. The simplest approach is an adaptation of Anderson's treatment of phase fluctuations and plasmons in a superconductor¹⁴ to the spin-triplet channel. Precisely this method, known as generalized RPA,^{15,16} was used by Demler *et al.*¹² To make the approach self-consistent, we make a quasiparticle approximation with a dispersion¹⁷ fitted to peak positions in the ARPES data in the superconducting state of $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$. As the measured dispersion is flat near $(0, \pi)$, our calculation yields a strikingly narrow two-electron continuum with a width of about 10 meV. This narrowness is caused by the proximity of the hot spots to the van Hove points at $(\pi, 0)$.

This quasiparticle model is a “best case” scenario for the antibound state. This model assumes that one has a superfluid Fermi liquid, which appears to be consistent with transport and ARPES data, at least at low temperatures. Within this approximation, the upper edge of the two-hole continuum is sharp, which is favorable for the formation of the π resonance. By making such

an approximation, we essentially neglect the large incoherent part of the electron spectral function known to exist from photoemission. On the other hand, this incoherent part is absent below the energy scale $\Delta + E_{\text{res}}$ set by the spin resonance,^{10,18} so the fermionic incoherence affects the two-particle propagators only at energies above $2\Delta + E_{\text{res}}$. Thus the π resonance can, in principle, live below this threshold. We make no attempt to explain *why* the quasiparticle dispersion is so remarkably flat near $(\pi, 0)$. A number of authors^{18–20} have pointed out that this flatness most likely results from a strongly ω -dependent self-energy which renormalizes the bare dispersion. Also note that the extreme narrowness of the two-particle continuum is not a crucial part of our argument: we have observed similar behavior with a much wider (70 meV) two-particle continuum based on normal state dispersions.

We now proceed with the calculations. Consider a system of fermions with dispersion $\epsilon_{\mathbf{k}}$ given in Ref. 17 and the nearest-neighbor interaction^{11,21}

$$H_1 = \sum_{\langle ij \rangle} J \left(\mathbf{S}_i \cdot \mathbf{S}_j - \frac{1}{4} n_i n_j \right) + V n_i n_j. \quad (1)$$

The J term gives rise to an attraction in the particle-particle d-wave singlet channel and in the particle-hole triplet channel, where it can induce a spin exciton. The V term accounts for repulsion in the triplet pairing channel and gives rise to the π resonance. In the conventional t - J model, $V = 0$. Following Refs. 11 and 21, we consider a more general interaction and treat V and J as independent parameters.

Although we are interested in spin susceptibility, in a superconducting ground state an operator of spin has the same quantum numbers as an operator creating a spin-triplet pair. We therefore analyze the linear response for the set of three operators¹²

$$S_+ = N^{-1/2} \sum_{\mathbf{k}} a_{\mathbf{k}\uparrow}^\dagger a_{\mathbf{k}+\mathbf{Q}\downarrow}, \quad (2)$$

$$\pi = N^{-1/2} \sum_{\mathbf{k}} g_{\mathbf{k}} a_{\mathbf{Q}-\mathbf{k}\downarrow} a_{\mathbf{k}\downarrow}, \quad (3)$$

$$\bar{\pi} = N^{-1/2} \sum_{\mathbf{k}} g_{\mathbf{k}} a_{\mathbf{Q}-\mathbf{k}\uparrow}^\dagger a_{\mathbf{k}\uparrow}^\dagger. \quad (4)$$

Here $a_{\mathbf{k},\sigma}$ is an electron annihilation operator and $g_{\mathbf{k}} = \cos(k_x a) - \cos(k_y a)$. Operators S_+ , π and $\bar{\pi}$ destroy a bosonic excitation with the same momentum $\mathbf{Q} = (\pi/a, \pi/a)$ and spin $S_z = -1$, but with different charges 0 and ± 2 , respectively.

As a warmup exercise, consider first a hypothetical (Fermi liquid) non-superconducting ground state. The triplet channels (2–4) are decoupled by virtue of a charge $U(1)$ symmetry. The Fourier transform of the bare pair susceptibility

$$\chi_\pi^0(t) = -i\theta(t) \langle [\pi(t), \pi^\dagger(0)] \rangle|_{V=J=0} \quad (5)$$

is shown in Fig. 2(a). As anticipated, the two-hole continuum ($\omega < 0$) is strikingly narrow: its upper edge is at $|E_{\text{max}}| \sim 10$ meV. The $\omega^{-1/2}$ divergence near E_{max} is a van Hove-type singularity associated with the $(\pi, 0)$ points. Observe that, at $|\omega| > |E_{\text{max}}|$, $\chi_\pi^0(\omega)$ is purely real. On the other hand, the bare spin susceptibility $\chi_S^0(\omega)$ [shown in the insert of Fig. 2(a)] is complex for all frequencies.

Within the generalized RPA approximation, the full susceptibilities are given by

$$\chi_\pi(\omega) = \frac{\chi_\pi^0(\omega)}{1 - V\chi_\pi^0(\omega)}, \quad \chi_S(\omega) = \frac{\chi_S^0(\omega)}{1 + 2J\chi_S^0(\omega)}. \quad (6)$$

Note the opposite signs of the interaction terms: attractive in the particle-hole channel, repulsive in the particle-particle channels. Because $\chi_\pi^0(\omega) = 1/V$ has a solution for any $V > 0$ [Fig. 2(a)], the full π susceptibility $\chi_\pi(\omega)$ acquires a pole above the upper edge of the continuum. This pole is the π resonance. In contrast, no pole occurs in the RPA response for S_+ . This is a consequence of the fact that the particle-hole continuum [and, hence, $\text{Im}\chi_S(\omega)$] extends to the lowest energies.

We now turn to the superconducting state. To facilitate the RPA treatment, we assume that the superconducting state is of the BCS type with a d -wave gap $\Delta(\mathbf{k}) = \Delta(\cos k_x a - \cos k_y a)/2$. We treat Δ as another input parameter in the problem. In principle, the gap has to be computed self-consistently, but for our purposes this is not necessary as we are not concerned with response functions in the singlet channel at small momenta.

Since in a superconductor charge is defined modulo 2, all three operators (2–4) now carry identical quantum numbers and one can use any superposition of these. It is customary to choose

$$A_0 = S_+, \quad A_1 = \frac{\pi - \bar{\pi}}{\sqrt{2}}, \quad A_2 = \frac{\pi + \bar{\pi}}{\sqrt{2}}. \quad (7)$$

The operators A_1 and A_2 describe fluctuations of the phase and the amplitude of the π mode, respectively.¹² The bare susceptibilities $\chi_{qq'}^0(\omega)$ are defined, similarly to Eq. (5), as the Fourier transforms of

$$\chi_{qq'}^0(t) = -i\theta(t) \langle [A_{q'}(t), A_q^\dagger(0)] \rangle|_{V=J=0}. \quad (8)$$

They can be written as components of a 3×3 matrix $\hat{\chi}^0$. At $T = 0$, we have

$$\chi_{qq'}^0(\omega) = \frac{1}{N} \sum_{\mathbf{k}} \left(\frac{\phi_q \phi_{q'}}{\omega - E_{\mathbf{k}} - E_{\mathbf{k}'}} - \frac{(-1)^{q+q'} \phi_q \phi_{q'}}{\omega + E_{\mathbf{k}} + E_{\mathbf{k}'}} \right), \quad (9)$$

where $E_{\mathbf{k}}^2 = \epsilon_{\mathbf{k}}^2 + \Delta_{\mathbf{k}}^2$, $\mathbf{k}' = \mathbf{Q} - \mathbf{k}$, and ϕ_q are components of the vector

$$\phi = \left(2^{-1/2} p(\mathbf{k}, \mathbf{k}'), \quad g_{\mathbf{k}} n(\mathbf{k}, \mathbf{k}'), \quad g_{\mathbf{k}} l(\mathbf{k}, \mathbf{k}') \right), \quad (10)$$

where l , n and p are the BCS coherence factors²²

$$\begin{aligned} l(\mathbf{k}, \mathbf{k}') &= u_{\mathbf{k}} u_{\mathbf{k}'} + v_{\mathbf{k}} v_{\mathbf{k}'}, \\ n(\mathbf{k}, \mathbf{k}') &= u_{\mathbf{k}} u_{\mathbf{k}'} - v_{\mathbf{k}} v_{\mathbf{k}'}, \\ p(\mathbf{k}, \mathbf{k}') &= u_{\mathbf{k}} v_{\mathbf{k}'} + v_{\mathbf{k}} u_{\mathbf{k}'}. \end{aligned} \quad (11)$$

The momentum sums are evaluated directly by including a small broadening $\omega \rightarrow \omega + i\Gamma$ in the energy denominators. Note that, due to strong particle-hole asymmetry implied by our dispersion, off-diagonal terms in $\hat{\chi}^0$ are by no means small.

Within the generalized RPA scheme, the full susceptibilities $\chi_{qq'}(\omega)$ are given by the matrix RPA equation

$$\hat{\chi}(\omega) = \hat{\chi}^0(\omega) + \hat{\chi}(\omega) \hat{V} \hat{\chi}^0(\omega). \quad (12)$$

where $\hat{V} = \text{diag}(-2J, V, V)$ in the basis (7). Recall that both J and V are assumed to be positive. We next discuss the behavior of response functions in various situations.

$J \neq 0$, $V = 0$. This is the situation considered in most studies.^{4,6,8,9} Although the BCS condensate now mixes all components of $\chi_{qq'}^0(\omega)$, the full spin susceptibility $\chi_{00}(\omega) = \chi_S(\omega)$ is still given by the standard RPA expression, Eq. (6).⁹ This effective decoupling is due to the fact that the J term in (1) yields no interaction in the particle-particle triplet channel. Despite a formal similarity with the normal state result, the form of $\chi_{00}^0(\omega)$ changes dramatically in a d-wave superconducting state: the Stoner continuum develops a hard gap

$$E_{\min}(\mathbf{Q}) = \min(E_{\mathbf{k}} + E_{\mathbf{Q}-\mathbf{k}}) = 2|\Delta(\mathbf{k}_{\text{hs}})| \approx 2\Delta. \quad (13)$$

The imaginary part of $\chi_{00}^0(\omega)$ jumps from 0 to a finite value at $2\Delta(\mathbf{k}_{\text{hs}})$. By Kramers-Kronig relation, this discontinuity causes a logarithmical divergence of $\text{Re}\chi_{00}^0(\omega)$. Therefore, for arbitrary J , the full $\chi_{00}(\omega)$ has a resonance pole at $\omega < 2\Delta$.⁶⁻¹⁰

In the absence of the triplet-pair coupling V , there is no π resonance above the continuum. At the same time, the π - π correlation function contains a resonance below the continuum, which is nothing but the spin exciton mixing into all other triplet channels.^{9,13} Indeed, for $V = 0$, the solution of Eq. (12) can be written in the following form:

$$\chi_{qq'}(\omega) = \chi_{qq'}^0(\omega) - 2\chi_{q0}^0(\omega) J^{\text{RPA}}(\omega) \chi_{0q'}^0(\omega), \quad (14)$$

where $J^{\text{RPA}}(\omega) = J/[1 + 2J\chi_{00}^0(\omega)]$. The spin-exciton pole in the RPA vertex $J^{\text{RPA}}(\omega)$ shows up in all full susceptibilities.

$J = 0$, $V \neq 0$. In this limit, there is no bound state in the particle-hole ($q = 0$) channel because $J = 0$. The π resonance in the particle-particle channel, which already existed in the normal state, is pushed to a higher energy when the superconducting gap opens up. We stress once again that the lower boundary of the two-hole continuum $E_{\min}$ is produced by fermions with momenta near hot spots, hence $E_{\min} \approx 2\Delta$. Certainly, an antibound state

in the particle-particle triplet channel is located above 2Δ . How much above 2Δ depends on the width of the two-electron continuum and the coupling strength.

The bare π - π response function in a superconducting state is presented in Fig. 2(b). It has two step-like discontinuities at $E_{\min}$ and $E_{\max}$, which are seen in both hole-hole ($\omega < 0$) and particle-particle ($\omega > 0$) spectra. The location of the π resonance in a superconductor is, however, not simply given by (6), but is a solution of the secular equation

$$\det \left[\begin{pmatrix} \chi_{11}^0(\omega) & \chi_{12}^0(\omega) \\ \chi_{21}^0(\omega) & \chi_{22}^0(\omega) \end{pmatrix} - \frac{1}{V} \right] = 0. \quad (15)$$

All four matrix elements $\chi_{qq'}^0$ for $q = 1, 2$ have the same thresholds as $\chi_{\pi\pi}^0$ in Fig 2(b). We solved this equation numerically and found that the π resonance indeed moves to an energy higher than 2Δ . For $V = 40$ meV, the resonance moves from 22 meV in the normal state [Fig. 2(a)] to 82 meV in the superconducting state with $2\Delta = 70$ meV [Fig. 2(b)]. Reasoning similar to that above implies that the π resonance shows up in the spin channel.

$J \neq 0$, $V \neq 0$. The general case interpolates between the two limits. Fig. 3 presents our results for the spin susceptibility at various ratios of the coupling strengths V/J . Interestingly enough, for comparable couplings, we found that the response functions contain *two* peaks at different energies. One, above $E_{\max}$, is the π resonance, the other, below $E_{\min}$, is the bound state in the particle-hole channel. As V is increased, the high-energy resonance pulls away from the upper edge of the two-hole continuum and strengthens. At the same time, the low-energy resonance approaches the lower edge and enters into the two-electron continuum, where it broadens and finally becomes invisible. In the opposite limit, as V gets progressively smaller, the π resonance weakens, merges with the outer edge of the continuum and disappears.

Discussion. Direct comparison²³ of neutron and ARPES data for $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ shows that the neutron resonance is located well below 2Δ . We believe that the same holds true for $\text{YBa}_2\text{Cu}_3\text{O}_{7-y}$. Because the lower continuum edge $E_{\min}$ is just below 2Δ , the resonance almost certainly occurs below this edge. This point can be verified by analysing ARPES data at the hot spots. In a slightly overdoped $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ sample with $T_c = 87$ K, the gap value at the hot spots is 32 ± 3 meV,²⁴ and $E_{\min} \geq 58$ meV. The neutron resonance at $\mathbf{Q} = (\pi/a, \pi/a)$ in the overdoped $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ with similar $T_c = 83$ K is observed at $E = 38$ meV,^{2,25} i.e., well below $E_{\min}$. We thus conclude that the neutron resonance is an exciton-like bound state in the particle-hole channel, not an antibound state, such as the π resonance.

It has been previously remarked that particle-hole mixing can mask the origin of the neutron resonance:^{9,12,13} a spin resonance mixes into triplet-pair channels and vice versa. We rely on a continuity argument to make the unambiguous identification of the experimental neutron peak, which is observed below the continuum edge

$E_{\min} \approx 2\Delta$, as expected for $V \ll J$. As the pair-triplet coupling V is turned off, the low-energy peak stays below the edge and continuously evolves into the usual RPA spin resonance at $V = 0$. On the other hand, there is no collective mode below $E_{\min}$ in the limit $V \gg J$, in which the π resonance can be defined unambiguously. Moving the high-energy π resonance across the continuum involves a discontinuous change. For this reason, the neutron resonance should not be associated with the π resonance.

We feel that a more realistic analysis will not change this general conclusion because our argument does not rely on detailed dynamics of the resonances, but is based on measured kinematics of low-energy fermion excitations and on general properties of bound states. If one wishes to find the π resonance in neutron scattering, the search should be confined to energies above the two-hole continuum. As its true upper edge is not known experimentally, we can only suggest that this energy exceeds 2Δ . On the other hand, because of the large incoherent parts of the spectral function observed in ARPES data for energies beyond Δ , there may be no true upper edge to the continuum,²¹ and it is quite possible that the π resonance will be strongly damped, or perhaps even absent, in neutron data.

The authors thank Ar. Abanov, J. C. Campuzano, E. Demler, B. Jankó, and J. F. Zasadzinski for helpful discussions. A. C. and O. T. are grateful to ANL staff for hospitality during their stay at Argonne, and M. N. to the staff at the ITP. This work was supported by the DOE Office of Science under contract W-31-109-ENG-38 (Argonne), by the DOE Grant No. DE-FG02-90ER40542 (Princeton) and by the NSF grants DMR-9979749 (Wisconsin) and PHY94-07194 (ITP).

¹ J. Rossat-Mignod, L. P. Regnault, C. Vettier, P. Bourges, P. Burllet, J. Bossy, J. Y. Henry, and G. Lapertot, *Physica C* **185-189**, 86 (1991).

² H. F. Fong, P. Bourges, Y. Sidis, L. P. Regnault, A. Ivanov, G. D. Gu, N. Koshizuka, and B. Keimer, *Nature* **398**, 588 (1999).

³ H. A. Mook, M. Yethiraj, G. Aeppli, T. E. Mason, and T. Armstrong, *Phys. Rev. Lett.* **70**, 3490 (1993).

⁴ M. Lavagna and G. Stemann, *Phys. Rev. B* **49**, 4235 (1994).

⁵ A. A. Abrikosov, *Phys. Rev. B* **57**, 8656 (1998).

⁶ D. Z. Liu, Y. Zha, and K. Levin, *Phys. Rev. Lett.* **75**, 4130 (1995).

⁷ I. I. Mazin and V. M. Yakovenko, *Phys. Rev. Lett.* **75**, 4134 (1995).

⁸ N. Bulut and D. J. Scalapino, *Phys. Rev. B* **53**, 5149 (1996).

⁹ J. Brinckmann and P. A. Lee, *Phys. Rev. Lett.* **82**, 2915

(1999); *J. Phys. Chem. Solids* **59**, 1811 (1998).

¹⁰ Ar. Abanov and A. V. Chubukov, *Phys. Rev. Lett.*, **83**, 1652 (1999).

¹¹ E. Demler and S.-C. Zhang, *Phys. Rev. Lett.* **75**, 4126 (1995).

¹² E. Demler, H. Kohno, and S.-C. Zhang, *Phys. Rev. B* **58**, 5719 (1998).

¹³ M. Greiter, *Phys. Rev. Lett.* **79**, 4898 (1997).

¹⁴ P. W. Anderson, *Phys. Rev.* **112**, 1900 (1958).

¹⁵ G. Rickayzen, *Phys. Rev.* **115**, 795 (1959).

¹⁶ A. Bardasis and J. R. Schrieffer, *Phys. Rev.* **121**, 1050 (1961).

¹⁷ M. R. Norman, *Phys. Rev. B* **61**, 14751 (2000). Note, dispersion “two” of Table I of that paper is used here.

¹⁸ M. R. Norman and H. Ding, *Phys. Rev. B* **57**, R11089 (1998).

¹⁹ E. Dagotto, A. Nazarenko, and M. Boninsegni, *Phys. Rev. Lett.* **73**, 728 (1994).

²⁰ A. J. Millis, *Phys. Rev. B* **45**, 13047 (1992); A. V. Chubukov, *Phys. Rev. B* **52**, R3840 (1995).

²¹ G. Baskaran and P. W. Anderson, *J. Phys. Chem. Solids* **59**, 1780 (1998).

²² J. R. Schrieffer, *Theory of Superconductivity* (Benjamin-Cummings, Reading, 1964).

²³ J. F. Zasadzinski, L. Ozyuzer, N. Miyakawa, K. E. Gray, D. G. Hinks, and C. Kendziora, preprint.

²⁴ H. Ding, M. R. Norman, J. C. Campuzano, M. Randeria, A. F. Bellman, T. Yokoya, T. Takahashi, T. Mochiku, and K. Kadowaki, *Phys. Rev. B* **54**, R9678 (1996).

²⁵ H. He, Y. Sidis, P. Bourges, G. D. Gu, A. Ivanov, N. Koshizuka, B. Liang, C. T. Lin, L. P. Regnault, E. Schönherr, and B. Keimer, cond-mat/0002013.




FIG. 1. “Hot spots” are points on the Fermi surface connected by the wave vector $\mathbf{Q} = (\pi/a, \pi/a)$ or those equivalent to it. Equivalently, they are given by intersections of the Fermi surface with the magnetic zone boundary (dashed lines).





FIG. 2. The bare response function ($T = 0$) of the operator π in a hypothetical Fermi liquid (a) and a d-wave superconductor with $\Delta_{\text{max}}=35$ meV (b) at $\mathbf{Q}$. Solid line: real part, dashed line: imaginary part. Insert: spin susceptibility in the normal state (note the difference in vertical scales). A broadening of $\Gamma=0.5$ meV was employed. Solution of the equation $\chi_{\pi}^0(\omega) = 1/V$ gives the RPA energy of the π resonance, $\omega = -E_{\pi}$. Note the difference in vertical scales. Horizontal arrows indicate two-particle continua.




FIG. 3. Spin susceptibility $[-\text{Im}\chi_S(\omega)]$ in the superconducting state at $\mathbf{Q}$. Values of V are shown near each curve. E_{min} and E_{max} are boundaries of the two-hole continuum. The chosen values for J and Δ are similar to those measured by neutron and ARPES, respectively.